

# BILOSIN 200

*Tylosin 200mg*

## INDICATIONS

Bilosin 200 is indicated for the treatment of diseases involving bacteria sensitive to Tylosin Including;

- Enzootic Pneumonia
- Swine Dysentery
- Erysipelas
- Respiratory diseases
- Secondary bacterial infection to virus disease
- Infectious Arthritis
- Cellulitis
- Otitis
- Mastitis
- Genito-Urinary infections



## BENEFITS

- Tylosin base injection
- Excellent syringability
- Effective against a wide range of Gram-negative and Gram-positive organisms

## PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1BIL001	100ml	12

Bimeda<sup>®</sup>  
 Broomhill Road  
 Tallaght  
 Dublin 24, Ireland  
 Tel: 1850-515253

See reverse side for full indications, administration and dosage.



[www.bimeda.ie](http://www.bimeda.ie)

## BILOSIN 200 INJECTION

### PRESENTATION

A clear yellow sterile solution suitable for parenteral administration. Each ml contains 200mg Tylosin Base BP(Vet). Also contains Benzyl Alcohol 4% as a preservative.

### USES

For the treatment of diseases in pigs involving organisms sensitive to Tylosin, including swine erysipelas (*Erysipelothrix rhusiopathiae*) vibriotic dysentery and pneumonia (*Mycoplasma hyopneumoniae*).

### DOSAGE AND ADMINISTRATION

0.5ml/10kg bodyweight, equivalent to 10mg of Tylosin per kg bodyweight, by intramuscular injection every 12 hours, up to a maximum of 6 injections. Do not inject more than 5ml at a single injection site. If a larger injection volume is necessary, it should be divided and administered at different injection sites.

### CONTRA-INDICATIONS, WARNINGS, ETC.

Do not use in animals known to be hypersensitive to the active ingredient.

Avoid skin contact with the preparation.  
Wash hands after use.

### WITHDRAWAL PERIODS

Pigs must not be slaughtered for human consumption during treatment. Pigs may be slaughtered for human consumption only after 21 days from the last treatment.

### PHARMACEUTICAL PRECAUTIONS

Do not store above 25°C.

Keep out of reach and sight of children.  
Following withdrawal of the first dose, use the product within 28 days.  
Avoid the introduction of contamination during use.  
Should any apparent growth or discoloration occur, the product should be discarded.

Legal Category POM

### PACKAGE QUANTITIES

Multi-dose vials of 100ml.

### VPA 10126/40/1

TECHNICAL INFORMATION  
Bimeda® a division of the Cross  
Vetpharm Group Ltd.  
Broomhill Road

